


# Dynamic Outcomes-Based Clustering of Disease Trajectory in Mechanically Ventilated Patients

Emma Rocheteau<sup>1</sup>, Dr Ioana Bica<sup>2</sup>, Prof. Pietro Liò<sup>1</sup>, and Dr Ari Ercole<sup>1</sup>

<sup>1</sup>University of Cambridge, UK

<sup>2</sup>University of Oxford, UK

# Mechanical Ventilation



# Why Cluster Patients?

- ▶ Patients on mechanical ventilation are highly heterogeneous.
- ▶ Clustering would help to generate:
  - ▶ Interpretable early warning systems.
  - ▶ Further understanding of disease trajectories.
  - ▶ Early categorisation of patients for intervention studies.


# Data: Electronic Health Records in Intensive Care

## AmsterdamUMCdb


- ▶ 14,836 ventilation episodes.
- ▶ Contains:
  - ▶ Vital Signs e.g. heart rate, blood pressure
  - ▶ Lab Results e.g. blood glucose
  - ▶ Demographics e.g. age, sex, ethnicity




# Methods




# Outcome Task Performance: Binary




# Outcome Task Performance: Duration




# Cluster Analysis




# Latent Space Visualisation




# Patient Trajectories



# The clusters are remarkably stable over time




# Most patients only appear in one cluster





# This happens very early in their admission






# Cluster Reliability


We tested:

- ▶ Alternative encoders
- ▶ Different initialisation seeds
- ▶ Varying the number of clusters


# Alternative Encoders



# Different Initialisation Seeds



# Number of Clusters



# “Stable” Cluster Transitions

Transition	Count	Median Time	Mortality (%)	Tracheostomy (%)	Urgency (%)	VD	LoS
3→1	17	3	76.5	0.0	47.1	0.5	0.7
5→1	29	16	51.7	10.3	55.2	4.3	5.3
1→3	28	11	10.7	0.0	67.9	1.0	2.6
5→3	46	9	15.2	4.3	41.3	1.2	6.5
2→4	28	17	10.7	21.4	42.9	6.2	12.8
5→4	27	10	11.1	7.4	48.1	3.4	9.1
1→5	25	3	44.0	4.0	68.0	3.9	6.5
3→5	15	4	13.3	13.3	53.3	1.9	4.6
4→5	15	56	26.7	26.7	46.7	6.6	11.5

# Summary

1. The TPC model outperforms alternative encoders.
2. We can generate clinically interpretable clusters.
3. The clusters are remarkably stable across time, and membership is determined early on.
4. Stable cluster transitions do occur, and are an important avenue for future work.

# Thank you!

## To my funders:

- ▶ The Armstrong Fund
- ▶ The Frank Elmore Fund
- ▶ The Cambridge University Clinical School

## ...and my wonderful coauthors:

Ioana Bica, Prof. Pietro Liò and Dr Ari Ercole

## ...and to W3PHIAI!

My email (Emma Rocheteau): [ecr38@cam.ac.uk](mailto:ecr38@cam.ac.uk)